





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07J 1/00	A1	(11) International Publication Number: WO 98/40398 (43) International Publication Date: 17 September 1998 (17.09.98)
<p>(21) International Application Number: PCT/US98/05026</p> <p>(22) International Filing Date: 13 March 1998 (13.03.98)</p> <p>(30) Priority Data: 08/816,558 13 March 1997 (13.03.97) US</p> <p>(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 08/816,558 (CON) Filed on 13 March 1997 (13.03.97)</p> <p>(71) Applicant (for all designated States except US): PHARM-ECO LABORATORIES, INC. [US/US]; 128 Spring Street, Lexington, MA 02173 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): SACHDEVA, Yesh [US/US]; 324 Hayward Hill Road, Concord, MA 01742 (US). RAM, Siya [US/US]; 9 Grant Road, Winchester, MA 01890 (US).</p> <p>(74) Agents: PIERCE, N., Scott et al.; Hamilton, Brook, Smith & Reynolds, P.C., Two Militia Drive, Lexington, MA 02173 (US).</p>	<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
<p>(54) Title: SYNTHESIS OF 2-ALKOXYESTRADIOLS</p> <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p>(a)</p> </div> <div style="text-align: center;">  <p>(b)</p> </div> </div> <p>(57) Abstract</p> <p>Disclosed is a method of preparing a compound represented by structural formula (a). The method comprises reacting bromine (Br₂) and an aliphatic organic acid with a compound represented by structural formula (b). R₁ and R₂ are each independently a hydroxyl protecting group.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

-1-

SYNTHESIS OF 2-ALKOXYESTRADIOLS

RELATED APPLICATION

This application is a continuation of U.S.S.N. 08/816,558 which was filed on March 13, 1997, the entire teachings of which are incorporated herein by reference.

GOVERNMENT FUNDING


This invention was made with government support under National Cancer Institute Contract No. NOI-CM-27764. The government has certain rights in the invention.

BACKGROUND OF THE INVENTION

At least some 2-alkoxy-estra-3,17 β -diols (hereinafter "2-alkoxy estradiols") have anticancer activity. For example, when tested in mice, 2-methoxy estradiol (1) acts as a potent inhibitor of neovascularization of solid tumors and inhibits their growth at doses which produce no apparent signs of toxicity (Fotsis et al., *Nature* 368:237 (1994)). Other 2-alkoxy estradiols that are believed to have anti-cancer activity are disclosed in co-pending U.S. provisional patent application "Novel 2-Alkoxy Estradiols and Derivatives Thereof", filed on March 13, 1997 (Attorney Docket No. PEL95-15p). 2-Methoxy estradiol (1) is thought to inhibit tumor growth act by:

- 1) inhibiting DNA synthesis and mitosis (Breuer et al., *Naturwissenschaft* 12:280 (1960) and Gelbke et al., *J. Steroid Biochem*, 7:457 (1976));
- 2) inhibiting tubulin polymerization or causing the formation of tubulin polymer with altered morphology and stability properties (D'Amato et al., *Proc. Natl. Acad. Sci. U.S.A.*


-2-




(1)

91:3964 (1994)); or 3) inhibiting angiogenesis, i.e. the creation of new blood vessels required for the growth of solid tumors (Fotsis et al., *Nature* 368:237 (1994), Folkman et al., *Nature* 339:58 (1994) and Blood et al., *Biochim. Biophys. Acts* et al., 1032:89 (1990)).

Syntheses previously employed to prepare 2-alkoxy estradiols have often suffered from low yields and a need to utilize tedious purification techniques. For example, one reported synthesis of 2-methoxy estradiol (1) involves the preparation of 2-formyl- β -estradiol dibenzyl ether (2)



(2)



(3)

-3-


as a key intermediate by formylation of β -estradiol (3), followed by benzylation of the two hydroxyl groups (Cushman et al., *J. Med. Chem.* 38:2041 (1995)). However, yields less than 10% are typically obtained from the two reactions. In addition, because the formylation reaction gives a mixture of regioisomers, 2-formyl- β -estradiol dibenzyl ether (2) must be purified by chromatography, which is generally considered to be an inefficient method of purification for large scale syntheses.

Therefore, a need exists for a method of synthesizing 2-alkoxy estradiols that overcomes or minimizes the above-mentioned problems.

SUMMARY OF THE INVENTION


The present invention is directed to an improved method for synthesizing 2-alkoxy estradiols and to specific 2-alkoxy estradiols. The invention is also directed to improved methods for preparing intermediates useful in the preparation of 2-alkoxy estradiols.

One embodiment of the present invention is a method of preparing a compound represented by the following structural formula:



The method comprises reacting bromine (Br_2) or iodine (I_2) and an aliphatic organic acid with a compound represented by the following structural formula:


5



R_1 and R_2 are each independently a hydroxyl protecting group. Preferably R_1 and R_2 are each a benzyl or substituted benzyl group, more preferably a benzyl group. R_3 is $-\text{Br}$ or $-\text{I}$.

10 The organic acid is preferably a C1 to C5 aliphatic organic acid, preferably a C1-C5 aliphatic carboxylic acid, more preferably acetic acid.


Yet another embodiment of the present invention is a method of preparing a compound represented by the following
15 structural formula:



-5-


R_1 and R_2 are as described above. R_4 is selected from the group consisting of -H, a lower alkyl group and a substituted lower alkyl group. R_4 is preferably -H.

The method comprises reacting an organolithium reagent
5 and a synthetic equivalent of R_4CO^* with a compound
represented by the following structural formula:



R_1 - R_4 are as described above. The organolithium reagent is preferably *n*-butyllithium. The synthetic equivalent of
10 R_4CO^* is preferably dimethylformamide.

Another embodiment of the present invention is a compound represented by the following structural formula:



R_{10} and R_{20} are independently a benzyl group or a substituted benzyl group. Preferably, R_{10} and R_{20} are each a benzyl group. R_{30} is -Br or -I.

The present invention has many advantages. For example, the bromination and the halogen-metal exchange reactions disclosed herein can be used to prepare 2-formyl- β -estradiol dibenzyl ether (2) in higher overall yields than known methods. In addition, the bromination introduces bromine predominantly at the two position of β -estradiol dibenzyl ether (5) to give a novel brominated estradiol dibenzyl ether intermediate (4), which can be substantially purified by crystallization. Thus, the disclosed method of preparing 2-formyl- β -estradiol dibenzyl ether (2) provides higher yields than methods previously used to prepare this compound with decreased formation of the 4-bromo by-product. As noted above, 2-formyl- β -estradiol dibenzyl ether (2) is an intermediate in the synthesis of 2-alkoxy- β -estradiols. Using the method of the present invention, 2-alkoxy estradiols can be synthesized more economically and in higher overall yield than was previously possible.

BRIEF DESCRIPTION OF THE DRAWING

The Figure is a schematic representation of the synthesis of 2-alkoxy estradiols.


25 DETAILED DESCRIPTION OF THE INVENTION

The features and details of the invention will now be more particularly described below and pointed out in the claims. It will be understood that the particular

-7-

embodiments of the invention are shown by way of illustration and not as limitations of the invention. The principal features of this invention can be employed in various embodiments without departing from the scope of the invention.

2-Alkoxy estradiols and a numbering system for identifying each carbon atom in the estradiol ring system are shown in the following structural formula:



10

R^5 is a lower alkyl group or a substituted lower alkyl group.

As used herein, suitable lower alkyl groups include C1-C6 straight or branched chain hydrocarbons which can optionally contain one or more double or triple bonds.


Suitable substituents on a lower alkyl group or phenyl group include one or more halogens (e.g., fluoro, chloro, bromo and iodo), nitro, nitrile, $-NH_2$, $-NH$ (lower alkyl), $-NH$ (substituted lower alkyl), $-N$ (lower alkyl)₂, $-N$ (substituted lower alkyl)₂, carbonyl groups, $-CONH_2$, $-CONH$ (lower alkyl), $-CONH$ (substituted lower alkyl), -

CON(lower alkyl)₂, -CON(substituted lower alkyl)₂, -CO₂H, -COO(lower alkyl) and -COO(substituted lower alkyl).

Halogenated alkyl groups can contain more than one kind of halogen. Examples of suitable lower alkyl or substituted lower alkyl groups include methyl, ethyl, n-propyl, iso-propyl, trifluoromethyl, trifluoroethyl, NO₂-CH₂-CH₂-, (CH₃)₂N-CH₂-CH₂-, -CH₂CH₂-CO-R', wherein R₁ is -H, lower alkyl, substituted lower alkyl, -OH, -O lower alkyl, -O(substituted lower alkyl), -NH₂, -NH(lower alkyl), -NH(substituted lower alkyl); -N(substituted lower alkyl)₂ and -N(lower alkyl)₂.

Using the bromination and metallation reactions of the present invention, 2-alkoxy estradiols can be prepared from estradiol (3). The Figure shows schematically the preparation of 2-ethoxy estradiol (10) and 2-(2',2',2'-trifluoroethoxy)estradiol (9) by this synthesis. Each of the reactions in this synthesis is described in greater detail below.

A first intermediate is prepared from estradiol (3) by protecting the two hydroxyl groups to form a compound represented by Structural Formula (I):



(I)

R_1 and R_2 are each independently a hydroxyl protecting group. Preferably R_1 and R_2 are each a benzyl or substituted benzyl group, more preferably a benzyl group.

5 As used herein, hydroxyl refers to alcohols and phenolic groups. A "suitable protecting group" is substantially inert with respect to the reagents used in the subsequent reactions in the disclosed synthesis of 2-alkoxy estradiols and does not cause, for example,
10 undesired side reactions. Hydroxyl protecting groups are well known in the art and are described in, for example, Chapters 2 and 3 of Greene and Wuts, "Protective Groups in Organic Synthesis", John Wiley & Sons (1991), the entire teachings of which are incorporated into this application
15 by reference. The skilled artisan can select, using no more than routine experimentation, suitable groups for use in the disclosed synthesis as well as conditions for applying and removing the hydroxyl protecting groups.


Preferred hydroxyl protecting groups include benzyl
20 groups and substituted benzyl groups. Examples of suitable substituents for benzyl protecting groups include, but are not limited to, one or more of the following groups: lower alkyl, fluoro, chloro, (lower alkyl)-O-, nitro, phenyl, di(lower alkyl)aminocarbonyl and silyl. Protecting a free
25 hydroxyl group with a benzyl group or substituted benzyl group is referred to herein as "benzylating" the hydroxyl group.

Estradiols can be benzylated by reacting at least one equivalent of a benzyl chloride, bromide or iodide and at
30 least one equivalent of a base per free hydroxyl group

(see, for example, Venuti et al., J. Med. Chem. 31:2132 (1988)). Suitable bases include hydride bases such as sodium hydride, potassium hydride and lithium hydride. Typically, a molar excess of benzyl halide and sodium
5 hydride are used with respect to each free hydroxyl group, for example up to a 1-3 fold excess of base and up to a 1-5 fold excess of benzyl halide. A catalytic amount of an iodide salt is preferably added to the reaction mixture, e.g., sodium iodide, potassium iodide or a
10 tetraalkylammonium iodide. Suitable solvents include apolar diprotic solvents such as dimethylformamide, dimethyl sulfoxide and hexamethylphosphoramide. Suitable temperatures for carrying out the reaction range from about 0°C to about 50°C, and preferably at room temperature. The
15 starting material and base are preferably mixed at 0°C.

Methods of protecting a hydroxyl group are described in Greene and Wuts, pages 53-71 and 156-159) and are encompassed within the scope of this invention. Specific conditions for benzylating estradiol (3) are provided in
20 Example 1.

The first intermediate is reacted with bromine (Br_2) or iodine (I_2) and an aliphatic organic acid in a suitable solvent to form a second intermediate represented by Structural Formula (II):



(II)

R₁ and R₂ are as described above. R₃ is -Br or -I.

Examples of suitable aliphatic organic acids include
5 Cl-C5 aliphatic carboxylic acids, sulfonic acids, sulfinic
acids and phosphoric acids. Aliphatic carboxylic acids are
preferred; acetic acid is particularly preferred.

The first intermediate is brominated (or iodinated)
with at least one molar equivalent of bromine (or I₂).
10 Typically an excess of bromine, for example up to about a
three fold excess, can be used. Preferably, about a 10% to
about a 20% excess of bromine is used. Suitable
concentrations of bromine in the reaction mixture range,
for example, from between about .01 M to about 5.0 M, and
15 preferably between about 0.10 M and about 0.40 M.

Suitable solvents for the bromination reaction
include, for example, etheral solvents in which the organic
acid and estradiol are soluble, for example,
tetrahydrofuran (THF), dioxane or glyme. THF is preferred.
20 Suitable amounts of aliphatic organic acid include a
volume/volume ratio of about 1:5 to about 5:1 with respect
to the solvent, preferably about 3:2 with respect to

-12-


tetrahydrofuran. The reaction can be run, for example, at temperatures between about -10°C and about 45°C .

Preferably, reagents are mixed under ice bath cooling, after which the reaction is allowed to warm to room

5 temperature.

Specific conditions for brominating a first intermediate, specifically estra-3,17 β -diol dibenzyl ether (5), are provided in Example 2. The bromination reaction using the conditions described above and in Example 2 forms
10 predominantly the 2-regioisomer over the 4-regioisomer. In addition, the brominated product can be substantially purified by crystallization from, for example, ethyl acetate.

The second intermediate is reacted with an
15 organolithium reagent and a synthetic equivalent of R_4CO^+ to form a third intermediate represented by Structural Formula (III):



(III)

R¹ and R² are as described above. R⁴ is selected from the group consisting of -H, lower alkyl and substituted lower alkyl.

Organolithium reagents are well known in the art and
5 are described in Wakefield, "Organolithium Methods"
Academic Press Limited, (1988), the teachings of which are
incorporated herein by reference. Organolithium reagents
are capable of reacting with aryl bromides and iodides such
as bromobenzene to effect a halogen/lithium exchange to
10 produce a lithiated intermediate, which can then be reacted
with a suitable electrophile. This type of reaction is
described in greater detail on pages 107-08 of Wakefield.
Examples of suitable organolithium reagents include n-
butyllithium, sec-butyllithium, t-butyllithium,
15 methyllithium and lithium metal.

"Synthetic equivalent" has the meaning commonly
associated with the term in the art of synthetic organic
chemistry (see, for example, Warren, "Designing Organic
Syntheses: A Programmed Introduction to the Synthon
20 Approach", John Wiley & Sons, 1978, page 8, and March,
"Advanced Organic Chemistry", John Wiley Sons, 1985, page
422, Corey, *Pure Appl. Chem.*, 14:19 (1967) pages 20-23.
"Synthetic equivalent" is understood to mean a reagent
which functions as the equivalent of a structural unit in
25 an organic molecule so that after one or more reactions
with a precursor, the reagent adds that structural unit to
the precursor. For example, dimethylformamide is a
synthetic equivalent of ¹CHO and, under suitable
conditions, reacts with nucleophiles, including lithiated
30 intermediates, to add a -CHO moiety to the precursor.


-14-

Thus, dimethylformamide achieves the same ultimate result as if a reagent which generated R_4CO^+ directly were used. Suitable synthetic equivalents of R_4CO^+ include, for example, $\text{R}_4\text{CO-N}(\text{lower alkyl})_2$ (e.g., dimethylformamide),
5 $\text{R}_4\text{CO-X}$, wherein X is a halide (e.g., acetyl chloride), $-\text{O}-\text{CO}-(\text{lower alkyl})$ (e.g., acetic anhydride) and $-\text{O}-(\text{lower alkyl})$ (e.g., ethyl acetate). Preferably, R_4 is $-\text{H}$ and the synthetic equivalent of R_4CO^+ is dimethylformamide.

To produce the third intermediate, the second
10 intermediate is typically reacted with, for example, between about 1.0 to about 5.0 equivalents, preferably between about 2.0 and about 3.0 equivalents, of an organolithium reagent, preferably *n*-butyllithium. Examples of suitable solvents include anhydrous ethereal solvents
15 (e.g., diethyl ether, dimethoxyethane, dioxane, tetrahydrofuran or diglyme) or anhydrous hydrocarbon solvents (e.g., benzene, pentane or hexane). THF is preferred. Suitable reaction temperatures range from, for example, between about -100°C to to about -60°C ,
20 preferably between about -75° to about -85°C . Suitable concentrations of the second intermediate in the reaction mixture range between about 0.005 M to about 1.0 M, preferably between about 0.050 M to about 0.10 M. The temperature is maintained until halogen metal exchange is
25 complete, from about ten minutes to about five hours, and typically about one to two hours, at which time an excess (e.g., from about 1.1 to about 10 fold, typically about 2.5 fold) of the synthetic equivalent of $-\text{R}_4\text{CO}^+$ with respect to organolithium reagent is added to the reaction mixture.
30 The synthetic equivalent can be added neat or as a solution

in an anhydrous etheral or hydrocarbon solvent. Typically, the reaction is warmed, for example, at about 0.5 to about 3.0 hours after addition of the synthetic equivalent to between about -30°C to about room temperature, preferably about -10°C to about 10°C, and allowed to stir until the reaction is complete. Specific conditions for converting a second intermediate, specifically 2-bromo-estra-3,17β-diol dibenzyl ether (4), to the third intermediate are provided in Example 3.

10 The third intermediate is reacted with an oxidizing agent capable of oxidizing an aryl alkyl ketone R-CO-Ar to the corresponding aryl alkonate R-CO-O-Ar to form a fourth intermediate represented by Structural Formula (IV):



15 (IV)

The reaction of ketones or aldehydes in the presence of suitable oxidizing agents to give esters, as in the transformation of the third intermediate to the fourth intermediate, is commonly known in the field of organic chemistry as a "Baeyer-Williger Oxidation". This reaction


20

is described, for example, in March, "Advanced Organic Chemistry", third edition, John Wiley & Sons, pages 990-91 (1985) and references cited therein, the teachings of which are incorporated herein by reference.

5 Suitable oxidizing agents for carrying out a Baeyer-Williger Oxidation include peracids such as *m*-chloroperbenzoic acid (hereinafter "mCPBA"), perbenzoic acid and peracetic acid. The reaction is carried out in the presence of an acid catalyst, for example, sulfonic
10 acids such as a methyl sulfonic acid, benzene sulfonic acid and *p*-toluene sulfonic acid. Suitable solvents include halogenated solvents such as methylene chloride, chloroform and dichloroethane.

 Suitable amounts of oxidizing agent include between
15 about 1.1 to about 2.0 equivalents with respect to the third intermediate. Suitable amounts of acid catalyst include between about 0.1% to about 10% by weight with respect to the oxidizing agent, preferably about 3-7%. Additional oxidizing agent and acid catalyst can be added
20 if the reaction does not go to completion. Suitable concentrations of the third intermediate in the reaction mixture include between about 0.005 M to about 5.0 M, preferably between about 0.05 M to about 0.5 M. The reaction is typically performed at room temperature, but
25 can also be carried out, for example, at temperatures ranging from about 0°C to the about 45°C. Specific conditions for carrying out a Baeyer-Williger Oxidation on a third intermediate, specifically 2-formyl-estra-3,β17-diol dibenzyl ether (2), are provided in Example 4.

The ester group of the fourth intermediate is cleaved to form a fifth intermediate represented by Structural Formula (V):



5

(V)

The cleavage of esters includes hydrolysis or alcoholysis of esters, which are well known in the field of chemistry. Conditions for the hydrolysis or alcoholysis of esters are described, for example, in Marvel, *Org. Syn., Coll.*, 3:495
10 (1995) and Marvel, *Org. Syn., Coll.* 2:416 (1943), the teachings of which are incorporated herein by reference. Specific conditions for hydrolyzing the ester of a fourth chemical intermediate are provided Example 4.


"Cleaving an ester" also refers to reacting an ester
15 R-CO-O-R' with a reducing agent suitable for reducing the ester to alcohols R-CH₂-OH and R'-OH. Suitable reducing reagents are well known in the art and include, for example, lithium aluminum hydride, lithium borohydride and diborane. Conditions suitable for suitable for reducing an
20 ester R-CO-O-R' to alcohols R-OH and R'-OH with an ester

reducing agent are also well known in the art and are disclosed, for example, in Brown et al., *Aldrichimica Acta* 12:3 (1979) and references cited therein, the teachings of which are incorporated herein by reference.

5 There are many other reactions known to the skilled artisan which are suitable for cleaving an alkyl aryl ester, such as is found in the compound represented by Structural Formula (IV) to give a compound represented by Structural Formula (V). It is to be understood that these
10 reactions are also encompassed by the term "cleaving the ester group".

The fifth intermediate is alkylated to form a sixth intermediate represented by Structural Formula (VI):

15



(VI)

R₁, R₂ and R₅ are as defined above. As used herein, "alkylation" refers to converting a phenolic group aryl-OH to the alkyl aryl ether aryl-O-R₃, wherein R₃ is as defined
20 above. The alkylation of phenols is well known in the

-19-

field of organic synthesis and can be carried out, for example, by the "Williamson Reaction". Conditions for carrying out the Williamson Reaction as well as other reactions suitable for alkylating phenolic groups are described in, for example, March, "Advanced Organic Chemistry", third edition, John Wiley & Sons, pages 342-43 (1985) and references cited therein, the teachings of which are incorporated herein by reference.

The alkylation can be carried out, for example, by reacting the fifth intermediate with an alkylating agent R_X , wherein X is a leaving group such as -Br, -I, mesylate, tosylate, brosylate or other suitable sulfonate ester, triflate or -O-SO-OCH₃. Examples include an alkyl bromide, an alkyl iodide (e.g., methyl iodide, ethyl iodide or 2,2,2-trifluoro-iodo-ethane) or a dialkyl sulfate (e.g., dimethyl sulfate) in the presence of a suitable base (e.g., potassium hydroxide, potassium carbonate or mercury (I) oxide). Suitable amounts of alkylating agent and base range, for example, from about 1.0 equivalent to about 10.0 equivalents and from about 1.0 to about 3.0 equivalents, respectively. Additional base and alkylating agent can be added if the reaction does not go to completion.


The fifth intermediate is reacted at concentrations, for example, ranging from about 0.01 M to about 5.0 M, preferably from about 0.1 M to about 0.5 M. Suitable solvents include halogenated solvents such as methylene chloride and aprotic polar solvents such as dimethylformamide. The reaction temperature ranges, for example, from about room temperature to about 150°C; when

aprotic polar solvents are used, the reaction temperature preferably ranges from about 110°C to about 130°C.

Specific conditions for alkylating a compound represented by Structural Formula (VI), specifically estradiol (6) and estradiol (8) in the Figure, are provided in Examples 5 and 6B respectively.

The protected hydroxyl groups at the three and seventeen positions in the sixth intermediate are deprotected to give a 2-alkoxy estradiol represented by

10 Structural Formula (VII):



(VII)

R_5 is as defined above.

As discussed earlier, suitable conditions for removing phenolic protecting groups are known in the art and are described in, for example, Greene and Wuts, "Protective Groups in Organic Synthesis", John Wiley & Sons (1991). When the hydroxyl protecting groups are benzylic or substituted benzylic groups, the deprotection reaction is referred to herein as a "debenzylation". Protected hydroxyl groups can be debenzylated by hydrogenolysis over

-21-

a palladium catalyst, for example $\text{Pd}(\text{OH})_2\text{-C}$ in an ethereal solvent (e.g. THF, diethyl ether) or an alcoholic solvent (methanol or ethanol). Suitable pressures include from about 10 to about 120 psi, preferably from about 40-60 psi.

5 The hydrogenation can be carried-out, for example, at temperatures from about 15°C to about 45°C , but preferably at room temperature, and at concentrations of the sixth intermediate ranging, for example, from about 0.01 M to about 5.0 M. The reaction is continued until complete, and

10 can be monitored by thin layer chromatography. Specific conditions for debenzylating a hydroxyl represented by Structural Formula (VI), specifically estradiols (7) and (8) in the Figure, are provided in Example 6. Other conditions for debenzylating a hydroxyl group are disclosed

15 in Chapters 2 and 3 of Wuts and Greene and are within the scope of the present invention.

In another embodiment of the present invention, the sixth intermediate represented by Structural Formula (VI), can be prepared directly from a compound represented by

20 Structural Formula (III). Specifically, bromide is nucleophilically displaced by an alkali metal alkoxide $\text{R}_5\text{O-Y}^+$, wherein Y^+ is an alkali metal cation, in the presence of a catalytic amount of copper (I) iodide. The reaction is typically carried out in an aprotic dipolar solvent such

25 dimethylformamide. The reaction nucleophilic displacement of halides from aryl halides with nucleophiles in the presence of copper (I) iodide is well known in the art as the "Ullman Reaction". Conditions for carrying out the Ullman Reaction are described in March, "Advanced Organic

30 Chemistry", third edition, John Wiley and Sons, pages 597-

598, references cited therein, and in Vyas and Shah, Org. Syn. Coll. Vol. 3, page 836, the teachings of which are incorporated by reference. Specific conditions for carrying out this reaction are described in Example 6A.

- 5 The invention is illustrated by the following examples which are not intended to be limiting in any way.

EXEMPLIFICATION

Melting points were determined in capillary tubes on a Mel-Temp apparatus and are uncorrected. Spectra were
10 obtained as follows: CI mass spectra on a Finnegan 4000 spectrometer; FAB mass spectra and EI mass spectra on a Kratos MS50 spectrometer; ¹H NMR spectra on a Bruker AC 300 spectrometer; IR spectra on a Beckman IR-33 spectrophotometer, or a Nicolet FT-IR Impact 410
15 spectrophotometer. Microanalyses were performed by MicroAtlantic Laboratories, Atlanta, and all values were within $\pm 0.4\%$ of the calculated compositions.

EXAMPLE 1 - 3,17 β -Dibenzyloxyestra-1,3,5(10)-Triene (5)

Sodium hydride (2.6 g, 65 mmol) was added in four
20 portions to a solution of estra-3,17 β -diol (3) (1, 5.0 g, 18 mmol) in anhydrous DMF (150 mL) under argon at 0°C, and the resulting mixture was stirred at 0°C for 15 min. Benzyl bromide (12.5 mL, 105 mmol) was added, followed by tetrabutylammonium iodide (0.50 g, 1.35 mmol). Stirring
25 was continued for 2 days at room temperature. The reaction mixture was cooled to 0°C and 50% aqueous ethanol (20 mL) was added slowly, followed by dropwise addition of 3 N HCl

(18mL) at 0°C to 5°C. The compound was extracted into ether (3 x 200 mL). The ether layer was washed with water (100 mL), brine (100 mL) and dried over anhydrous sodium sulfate. Evaporation of the ether layer under reduced
5 pressure gave a viscous oil. Trituration with hexane afforded a white solid (5.0 g, 60%), which was collected by filtration and dried under vacuum: mp 74-78°C [Fuji et al., *J. Org. Chem.*, 44:1661-1664 (1979)., mp 81-82°C (ethanol, acetone); Hamacher, H.; Christ, E. Potential
10 Antineoplastics. 7th Comm: Introduction of a Nitrogen Mustard Group into the 6 α -Position of Estradiol. *Arzneim. Forsch.*, 33:347-352 (1983). mp 62°C (aq. Methanol)]; R_f 0.334 (hexane-ethyl acetate (95:5), silica gel]; IR (KBr) 2910, 2850, 1600, 745, 685 cm^{-1} ; ^1H NMR (CDCl_3) δ 0.9 (s, 3
15 H), 1.05-2.45 (m, 14H), 2.83 (m, 1 H), 3.5 (t, J = 8.5 Hz, 1 H), 4.57 (s, 2 H), 5.05 (s, 2 H), 6.67-6.82 (m, 2 H), 7.12-7.57 (m, 11 H). Anal. ($\text{C}_{32}\text{H}_{36}\text{O}_2$) C, H.

EXAMPLE 2 - 2-Bromo-3,17 β -Dibenzyloxyestra-1,3,5(10)-
Triene (4)

20 Bromine (0.58 mL, 11.3 mmol) was added by syringe to an ice-cold, stirred solution of estradiol (5) (4.5 g, 10 mmol) in an acetic acid/THF mixture (3:2, 45 mL). The resulting reaction mixture was allowed to warm to room temperature with stirring. After 1.5 h, the mixture was
25 poured onto an ice-water mixture (200 mL), and compound was extracted with dichloromethane (3 x 200 mL). The combined organic layer was washed with water (200 mL), saturated aqueous sodium bicarbonate (200 mL), 10% aqueous sodium thiosulfate (150 mL), water (100 mL), brine (100 mL), and

dried over anhydrous sodium sulfate. The organic layer, on evaporation under vacuum, followed by trituration with hexane-ethyl acetate mixture, gave the product (3.9 g, 75%). Only about 10% of the 4-bromo regioisomer is formed.

5 An analytical sample was prepared by crystallization from ethyl acetate; mp 156-158°C; IR (KBr) 2960-2840, 1600, 1590, 730-720, 685 cm^{-1} ; ^1H NMR (CDCl_3) δ 0.9 (s, 3H), 1.10-2.40 (m, 13H), 2.75-2.85 (m, 2H), 3.5 (t, $J = 8.5$ Hz, 1H), 4.6 (s, 2 H), 5.63 (s, 2 H), 6.67 (s, 1 H), 7.25-7.60 (m,
10 11 H). Anal. ($\text{C}_{32}\text{H}_{35}\text{BrO}_2$) C, H.

These results show that 2-bromo- β -estradiol dibenzyl ether (4) can be obtained in 75% yield from β -estradiol dibenzyl ether (5) with diminished formation of the 4-bromo by-product.

15

EXAMPLE 3 - 2-Formyl-3,17 β -Dibenzyloxyestra-1,3,5(10)-
Triene (2)

A 1.6 M solution of $n\text{-BuLi}$ (17.5 mL) in hexanes was added dropwise by syringe under argon atmosphere at -78°C
20 to a solution of estradiol (4) (5.88 g, 11.2 mmol) in anhydrous THF (150 mL), and the resulting reaction mixture was stirred at -78°C for 2 h. Anhydrous DMF (5.5 mL) was added and the stirring was continued for 1 h at -78°C . The mixture was warmed to 0°C and stirred for 1 h, and poured
25 onto an ice-cold solution of 3N HCl (100 mL). The aqueous layer was extracted with ether (3 x 300 mL). The combined organic layer was washed with 50% brine-water (100 mL), brine (150 mL), and dried over anhydrous sodium sulfate. The organic layer was filtered and evaporated under reduced
30 pressure to afford impure estradiol (2), which was purified

over a silica gel column using hexane-ethyl acetate as eluant. Appropriate fractions were combined and evaporated under reduced pressure to afford estradiol (2) (4.36 g, 82%): mp 145-147°C; IR (KBr) 2920, 2820, 1675, 1600, 728, 685 cm⁻¹, ¹H NMR (CDCl₃) δ 0.88 (s, 3 H), 1.1-2.5 (m, 13 H), 2.80-2.97 (m, 2 H), 3.50 (t, *J* = 8.5 Hz, 1 H), 4.56 (s, 2 H), 5.65 (s, 2 H), 6.75 (s, 1H), 7.20-7.50 (m, 10 H), 7.78 (s, 1 H), 10.5 (s, 1 H). Anal. (C₃₄H₄₀O₃)C, H.

These results show that 3,17-diprotected 2-bromo-β-estradiols can be converted in high yield to 3,17-diprotected 2-formyl-β-estradiols by a halogen-metal exchange followed by quenching with dimethylformamide.

EXAMPLE 4 - 2-Hydroxy-3,17β-Dibenzyloxyestra-1,3,5(10)-Triene (6)

mCPBA (3.3 g, 19.1 mmol) and *p*-toluenesulfonic acid monohydrate (0.16g, 0.84 mmol) were added successively to a solution of estradiol (2) (5.8g, 12.1 mmol) in anhydrous dichloromethane (150 mL). The resulting mixture was stirred at room temperature, and the reaction was followed by TLC (dichloromethane/silica gel). After 10 h, additional amounts of mCPBA (1.0 g) and *p*-toluenesulfonic acid (0.040 g) were added, and stirring was continued for 5 h. The reaction mixture was diluted with dichloromethane (300 mL), and the organic layer was washed with 10% sodium sulfite solution (100 mL), water (100 mL), brine (100 mL), and dried over anhydrous sodium sulfate. The organic layer on evaporation under reduced pressure afforded 4.85 g of crude 2-formyloxy-estra-3,17β-diol dibenzyl ether (4.85 g). This was suspended in anhydrous methanol (250 mL), four

-26-

drops of conc H_2SO_4 were added, and the resulting mixture was stirred at 90°C for 1h. The methanol was removed under reduced pressure at 50°C and the resulting residue was dissolved in dichloromethane (400 mL). The organic layer was washed with water (100 mL), saturated sodium bicarbonate solution (100 mL), water (100 mL), and brine (100 mL), and dried over anhydrous sodium sulfate. The dichloromethane layer on evaporation under reduced pressure gave crude compound (6) (4.15 g) as a gummy solid, which on purification over a silica gel column using hexane-dichloromethane mixture as an eluant provided pure estradiol (6) (3.8 g, 67%) as a white colored solid: mp $117-118^\circ\text{C}$; R_f 0.364 (dichloromethane, silica gel); IR (KBr) $3540, 2960-2840, 1510, 1450, 735-720, 685\text{ cm}^{-1}$; $^1\text{H NMR}$ (CDCl_3) δ 0.9 (s, 3 H), 1.05-2.40 (m, 14 H), 2.80 (m, 1 H), 3.50 (t, $J = 8.5\text{ Hz}$, 1 H), 4.60 (s, 2 H), 5.08 (s, 2 H), 5.5 (s, 1 H), 6.65 (s, 1 H), 6.82 (s, 1 H), 7.20-7.50 (m, 10 H). Anal. ($\text{C}_{32}\text{H}_{36}\text{O}_3$) C, H.

EXAMPLE 5 - 2-(2',2',2'-Trifluoroethoxy)-3,17 β -
Dibenzyloxyestra-1,3,5(10)-Triene (7)

Powdered K_2CO_3 (1.5 g, 11 mmol) was added to a solution of estradiol (6) (4.0 g, 8.5 mmol) in anhydrous DMF (50 mL) followed by dropwise addition of $\text{CF}_3\text{CH}_2\text{I}$ (5.0 mL, 51 mmol) at room temperature. The resulting reaction mixture was heated at 110°C for 3 h. Additional K_2CO_3 (2.5 g) and $\text{CF}_3\text{CH}_2\text{I}$ (6 mL) were added, and the mixture was heated again at 130°C for 2 h. The mixture was cooled in an ice bath and poured onto ice cold 3N HCl (125 mL). The aqueous layer was extracted with ether (2 x 200 mL). The combined

-27-

ether layer was washed with 3 N HCl (100 mL), water (100 mL), brine (100 mL), and dried over anhydrous MgSO_4 . The ether layer was evaporated under reduced pressure and the resulting crude product on purification on a silica gel column gave estradiol (7) (4.5 g, 95%) as a colorless oil:

5 IR (neat) 2910, 2850, 1600, 1500, 730, 690 cm^{-1} ; ^1H NMR (CDCl_3) δ 0.9 (s, 3 H), 1.1-2.4 (m, 13 H), 2.65-2.85 (m, 2 H), 3.50 (t, $J = 8.5$ Hz, 1H), 4.40 (q, $J = 8.5$ Hz, 2 H), 4.56 (s, 2 H), 5.05 (s, 2 H), 6.70 (s, 1 H), 6.98 (s, 1 H),

10 7.2-7.55 (m, 10 H). Anal. ($\text{C}_{26}\text{H}_{38}\text{O}_2$) C, H.

EXAMPLE 6 - 2-(2',2',2'-Trifluoroethoxy)estra-1,3,5(10)-Trien-3,17 β -Diol (9)

$\text{Pd}(\text{OH})_2\text{-C}(20\%, 1.0 \text{ g})$ was added carefully under argon atmosphere to a solution of estradiol (7) (1.0 g, 1.8 mmol) in anhydrous THF (50 mL). The resulting mixture was

15 hydrogenated at 45-50 psi on a Parr apparatus for 24 h. The catalyst was removed by filtration using a celite pad under argon atmosphere, and the pad was washed with dichloromethane (200 mL). Evaporation of the filtrate

20 under reduced pressure, followed by purification on a silica gel column, gave compound (9) (0.5 g, 75%): mp 167-168°C; IR (KBr) 3550, 2960-2840, 1590, 1510, 870 cm^{-1} ; ^1H NMR (CDCl_3) δ 0.80 (s, 3 H), 1.10-2.40 (m, 13 H), 2.70-2.85 (m, 2 H), 3.75 (m, 1 H), 4.40 (q, $J = 8.5$ Hz, 2 H), 5.35

25 (s, 1 H, OH), 5.5 (s, 1 H), 6.70 (s, 1 H), 6.80 (s, 1 H); MS m/z (relative intensity) 370 (100), 311 (9), 270 (8), 244 (8), 205 (6). Anal. ($\text{C}_{26}\text{H}_{38}\text{O}_3 \cdot 0.7 \text{ H}_2\text{O}$) C, H.

EXAMPLE 6A - 2-Ethoxy-3,17 β -dibenzyloxyestra-1,3,5(10)-triene(8):

To a stirred solution of sodium ethoxide (10.5 g, 154 mmol) in anhydrous ethanol (30 mL), anhydrous DMF was added under argon atmosphere and stirring was continued for 20 minutes at room temperature. To the above solution CuI (10.6 g, 56 mmol) and 2-bromocompound (4) (5.3 g, 10.0 mmol) were added successively, and the resulting mixture was stirred at 105°C to 110°C for 7 to 8 hours. The reaction was cooled and poured onto ice-cold water (200 mL), and followed by addition of 3 N HCl (25 mL). The insoluble material was filtered off through a celite pad and the aqueous layer was extracted with dichloromethane (300 mL x 3). The combined organic layer was washed with water (200 mL), brine (200 mL), and dried over sodium sulfate. The organic layer was filtered and evaporated under reduced pressure to afford a mixture which was purified over a silica gel column using hexane:ethyl acetate as an eluent. Appropriate fractions were combined and evaporated under reduced pressure. Yield of compound (8) was 60%; mp 76°C-77°C; IR (KBr) 2910, 2850, 740, 685 cm⁻¹. ¹H NMR (CDCl₃) δ 0.9 (s, 3H, -CH₃), 1.1-2.35 (m, 18H, -CH, -CH₂ & CH₃), 3.5 (t, 1H, -OCH), 4.10 (q, 2H, -OCH₂), 4.6 (s, 2H, -OCH₂), 5.60 (s, 2H, -OCH₂), 6.64 (s, 1H, Ar-H), 6.89 (s, 1H, Ar-H), 7.15-7.55 (m, 10H, Ar-H). Analysis calcd for C₃₄H₄₀O₃: C, 82.2; H, 8.12. Found: C, 82.51; H, 8.11. In addition to desired product (8), two minor by products 2-methoxy-3,17 β -dibenzloxyestra-1,3,5(10)-triene and 2-N,N-dimethylamino-3,17 β -dibenzyloxyestra-1,3,5(10)-triene were also obtained in 5% and 0.72% yield respectively.

-29-

EXAMPLE 6B - 2-Ethoxyestra-1,3,5(10)-trien-3,17 β -diol(10)

To a solution of 2-ethoxyderivative (8) (1.1 g, 2.3 mmol) in anhydrous THF (40 mL), 20% Pd (OH)₂-C(2.3 g) was added carefully under argon atmosphere. The resulting mixture was hydrogenated at 45-50 psi on a Parr Apparatus for four hours. The resulting mixture was monitored by TLC. After completion, the catalyst was removed via filtration using a celite pad under argon atmosphere, and pad was washed with dichloromethane (100 mL). Filtrate on evaporation under reduced pressure gave 0.8 g of impure compound, which was purified by column chromatography. Yield >95%; mp 154°C-156°C [lit. 154°C-156°C]; [α]_D +92.24° (c=0.508, CHCl₃) {lit. [α]_D = +90° (c=0.5, CHCl₃)}. IR (KBr) 3720, 3380, 2980-2840, 1580, 1510 cm⁻¹. ¹H NMR (CDCl₃) δ 0.83 (s, 3H, -CH₃), 1.10-2.38 (m, 16H, -CH, CH₂ & -CH₂), 2.70-2.90 (m, 2H, -CH₂), 3.75 (m, 1H, -OCH), 4.10 (q, 2H, -OCH₂), 5.5 (s, 1H, OH), 6.67 (s, 1H, Ar-H), 6.82 (s, 1H, Ar-H). Analysis calcd. For C₂₀H₂₈O₃. C, 75.91; H, 8.92. Found: C, 75.97; H, 8.96.

20 EXAMPLE 7 - Tubulin Assays

Electrophoretically homogeneous tubulin was purified from bovine brain as described previously (Hamel et al., *Biochemistry* 23:4173-4184 (1984)). Determination of IC₅₀ values for the polymerization of purified tubulin was performed as described in D'Amato et al. *Proc. Natl. Acad. Sci. U.S.A.* 91:3964-3968 (1994). Tubulin was preincubated at 26°C with varying compound concentrations, reaction mixtures were chilled on ice, GTP (required for the polymerization reaction) was added, and polymerization was

followed at 26°C by turbidimetry at 350 nm in Gilford recording spectrophotometers equipped with electronic temperature controllers. Four instruments were used, and two control reaction mixtures were present in each experiment. The extent of polymerization after a 20 min incubation was determined (the values for the two controls were usually within 5% of each other). IC_{50} values were determined graphically. Active compounds were examined in at least three independent assays, while inactive compounds (defined as IC_{50} value $> 40 \mu M$) were examined in at least two independent experiments. Inhibition of colchicine binding to tubulin was performed as described previously (D. Amato et al., *Proc. Natl. Acad. Sci. U.S.A.* 91:3964-3968 (1994)). Reaction mixtures contained 1.0 μM tubulin (0.1 mg/mL), 5.0 μM [3H]colchicine, and 50 μM inhibitor. Incubation was for 30 min at 37°C.

The inhibitory activities of 2-methoxy estradiol (1), 2-ethoxy estradiol (10), and 2-(2',2',2'-trifluoroethoxy) estradiol (9) on tubulin polymerization are reported in Table 1. Inhibition of tubulin polymerization was investigated using electrophoretically homogeneous tubulin from bovine brain. 2-ethoxy estradiol (10) was over three times more active than 2-methoxy estradiol (1).

-31-

EXAMPLE 8 - Activity of 2-alkoxy Estradiols and Analogs
Thereof in the National Cancer Institute
Anticancer Screen

The effects of the analogs on cancer cell growth
5 was investigated in the National Cancer Institute (NCI)
Developmental Therapeutics Program's *in vitro* panel of
approximately 55 cell lines (Grever et al., *Seminars in
Oncology* 19:622 (1992), Alley et al., *Cancer Research*
48:589 (1988) and Montes et al., *J. National Cancer*
10 *Institute* 83:757 (1991)). Representative results are
listed in Table 1 for HOP-62 non-small-cell lung cancer
cells, HCT-116 colon cancer cells, SF-539 central nervous
system (CNS) cancer cells, UACC-62 melanoma cells, OVCAR
ovarian cancer cells, SN12-C renal cancer cells, DU-145
15 prostate cancer cells, and MDA-MB-435 breast cancer cells.
2-Ethoxy estradiol (10) was more active against all the
cell lines listed in Table 1 than 2-methoxy estradiol (1).

**Table 1. Cytotoxicities and Antitubulin Activities of
2-Methoxyestradiol and Analog cytotoxicity (GI_{50} in μM)^a**

No.	Lung HOP-62	Colon HCT-116	CNS SF-539	Melanoma UACC-62	Ovarian OVCAR	Renal SN12-C	Prostate DU-145	Breast MDA-MB- 435	MGM ^b	Inhibn of Tubulin Polymn IC_{50} ($IC_{50}/\mu M$)
1	0.70	0.47	0.32	0.36	0.21	0.95	1.8	0.080	1.30	2.9 ± 0.6
5	11.6	3.22	1.38	2.32	0.46	4.46	7.05	0.28	2.63	1.7
14	0.018	0.026	0.014	0.016	0.016	0.039	0.065	<0.01	0.076	0.91 ± 0.5

^aThe cytotoxicity GI_{50} values are the concentrations corresponding to 50% growth inhibition.

^bMean graph midpoint for all human cancer cell lines (approximately 55) tested.

Equivalents


Those skilled in the art will be able to recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific
5 embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

CLAIMS


What is claimed is:

1. A method of preparing a compound represented by the following structural formula:

5



comprising reacting Br_2 or I_2 and an aliphatic organic acid with a precursor compound represented by the following structural formula:




-35-

wherein R_1 and R_2 are each independently a hydroxyl protecting group; and


R_3 is -Br or -I.

2. The method of Claim 2, wherein the precursor compound
5 is reacted with Br_2 and an aliphatic organic acid and
 R_3 is -Br.
3. The method of Claim 2 wherein:
 - a) the aliphatic organic acid is a C1-C5 aliphatic
carboxylic acid; and
 - 10 b) R_1 and R_2 are each a benzyl or substituted benzyl
group.
4. The method of Claim 3 wherein the aliphatic organic
acid is acetic acid and the hydroxyl protecting group
is a benzyl group.
- 15 5. The method of Claim 3 wherein the precursor compound
is prepared by benzylating a compound represented by
the following structural formula:




6. A method of preparing an estradiol represented by the following structural formula:

5



wherein R₅ is a lower alkyl group or a substituted lower alkyl group, from a starting compound represented by the following structural formula:


10



said method comprising the steps of:


- a) benzylating the starting compound to form a first intermediate represented by the following structural formula:

5




wherein R_1 and R_2 are each independently a benzyl or substituted benzyl group;


- b) reacting the first intermediate with Br_2 and a Cl-C5 aliphatic carboxylic acid to form a second intermediate represented by the following structural formula:



- c) reacting the second intermediate with an organolithium reagent and a synthetic equivalent of R_4CO^* , wherein R_4 is selected from the group consisting of -H, a lower alkyl group and substituted lower alkyl group, to form a third intermediate represented by the following structural formula:




- d) reacting the third intermediate with an oxidizing agent capable of oxidizing an aryl alkyl ketone to the corresponding aryl alkonate, to form a fourth intermediate represented by the following structural formula:




-40-

- e) cleaving the aryl alkanoate group of the fourth intermediate to form a fifth intermediate represented by the following structural formula:




- f) alkylating the free phenol group of the fifth intermediate with an alkylating agent R_5X , wherein X is a leaving group, to form a sixth intermediate represented by the following structural formula:



- g) debenzylating the sixth intermediate to form the estradiol.

15


7. The method of Claim 6 wherein:
- a) the C1-C5 aliphatic carboxylic acid is acetic acid;
 - b) the organolithium reagent is *n*-butyllithium;
 - 5 c) the synthetic equivalent of R_4CO^+ is dimethylformamide;
 - d) R_4 is -H; and
 - e) the oxidizing agent is *m*-chloroperbenzoic acid.
8. The method of Claim 7 wherein R_5 is selected from the
- 10 group consisting of methyl, ethyl, $(CH_2)_N-CH_2-CH_2-$, $NO_2-CH_2-CH_2-$ and 2,2,2-trifluoroethyl.
9. A method of preparing an estradiol represented by the following structural formula:



15

wherein R_5 is a lower alkyl group or a substituted lower alkyl group, from a starting compound represented by the following structural formula:


-42-



said method comprising the steps of:

- a) benzylating the starting compound to form a first intermediate represented by the following structural formula:

5




wherein R_1 and R_2 are each independently a benzyl or substituted benzyl group;


- b) reacting Br_2 and an aliphatic organic acid with the first intermediate to form a second intermediate represented by the following structural formula:

10

-43-



- b) reacting the second intermediate with an alkali metal alkoxide $R_5O^- Y^+$, wherein Y^+ is an alkali metal cation, and copper (I) iodide to give a third intermediate represented by the following structural formula:




- c) debenzylating the third intermediate to give the estradiol.

10. The method of Claim 9 wherein:

-44-

- a) R_1 and R_2 are benzyl groups;
b) R_3 is selected from the group consisting of methyl, ethyl, $(CH_3)_2N-CH_2-CH_2-O-$, $NO_2-CH_2-CH_2-O-$ and 2,2,2-trifluoroethyl; and
5 c) the aliphatic organic acid is acetic acid.


11. A method of preparing a compound represented by the following structural formula:



10 wherein:


R_1 and R_2 are each independently a phenol protecting group; and

R_4 is selected from the group consisting of -H, a lower alkyl group and a substituted lower alkyl group;
15 from a starting compound represented by the following structural formula:





comprising reacting the starting compound with an organolithium reagent and a synthetic equivalent of R_4CO^+ .

- 5 12. The method of Claim 11 wherein:
- a) R_1 and R_2 are each a benzyl group or a substituted benzyl group;
 - b) the organolithium reagent is *n*-butyllithium;
 - c) the synthetic equivalent of R_4CO^+ is
 - 10 dimethylformamide; and
 - d) R_4 is -H.
13. The method of Claim 12 wherein the starting compound
- is prepared by reacting Br_2 and an aliphatic organic acid with a compound represented by the following
- 15 structural formula:



14. The method of Claims 13 wherein the aliphatic organic acid is a C1-C5 aliphatic carboxylic acid and R_1 and R_2 are each a benzyl group.
- 5 15. The method of Claim 14 wherein the aliphatic organic acid is acetic acid.
16. A compound represented by the following structural formula:





INTERNATIONAL SEARCH REPORT

I. national Application No

PCT/US 98/05026

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07J1/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07J

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	C. AKIRA HORIUCHI ET AL: "Novel Regioselective Iodination of Estradiol 17.beta.-Acetate" BULLETIN OF THE CHEMICAL SOCIETY OF JAPAN., vol. 59, no. 8, 1986, TOKYO JP, pages 2459-2462, XP002067992 see page 2460, column 1, paragraph 3 see page 2461, column 2, paragraph 4 --- -/--	1



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

² Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

15 June 1998

Date of mailing of the international search report

09/07/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Watchorn, P

INTERNATIONAL SEARCH REPORT

National Application No

PCT/US 98/05026

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	XU-HUA ZENG ET AL: "A New Synthetic Route to Pyrogallolestrogen Dimethyl Ethers by Nucleophilic Substitution of 2,4-Dibromoestrogens" STERIODS., vol. 40, no. 2, August 1982, SAN FRANCISCO US, pages 121-124, XP002067993 see page 122, paragraph 2 ---	1-17
Y	CHEN S H ET AL: "A new synthetic route to 2- and 4-methoxyestradiols by nucleophilic substitution" STERIODS: STRUCTURE, FUNCTION, AND REGULATION., vol. 47, no. 1, January 1986, MA US, pages 63-66, XP002067994 see page 64 ---	1-17
Y	NAMBARA T; HONMA S; AKIYAMA S: " Analytical chemical studies on steroids. XXXIII. Steroid conjugates. III. New syntheses of 2-methoxyestrogens" CHEMICAL AND PHARMACEUTICAL BULLETIN., vol. 18, no. 3, 1970, TOKYO JP, pages 474-480, XP002067995 see the whole document ---	6-8
Y	HE H M ET AL: "A Novel 1,3 O.fwdarw. C Silyl Shift and Deacylation Reaction Mediated by Tetra-n-butylammonium Fluoride in an Aromatic System" JOURNAL OF ORGANIC CHEMISTRY., vol. 60, no. 18, 8 September 1995, EASTON US, pages 5905-5909, XP002067996 see scheme 1 see page 5906, column 1 ---	6-8, 14, 15
X	see page 5906, column 1 ---	11, 12
Y	CUSHMAN M ET AL: "SYNTHESIS, ANTITUBULIN AND ANTOMITOTIC ACTIVITY, AND CYTOTOXICITY OF 2-METHOXYESTRADIOL, AN ENDOGENOUS MAMMALIAN METALBOLIC OF ESTRADIOL THAT INHIBITS TUBULIN POLYMERIZATION BY BINDING TO THE COLCHICINE BINDING SITE" JOURNAL OF MEDICINAL CHEMISTRY, vol. 38, no. 12, 9 June 1995, pages 2041-2049, XP002059349 see page 2042, column 1, scheme 1; see page 2042, column 2, scheme 3 ---	6-8
	-/--	

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/05026

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	NUMAZAWA M; OGURA Y; KIMURA K; NAGAOKA M: "Synthesis of 2-methoxy- and 4-methoxyestrogens with halogen-methoxy exchange reactions" J. CHEM. RES. MINIPRINT, no. 11, 1985, pages 3701-3715, XP002067997 see the whole document -----	9, 10
P, X	CUSHMAN M ET AL: "Synthesis of Analogs of 2-Methoxyestradiol with Enhanced Inhibitory Effects on Tubulin Polymerization and Cancer Cell Growth" JOURNAL OF MEDICINAL CHEMISTRY., vol. 40, no. 15, 18 July 1997, WASHINGTON US, pages 2323-2334, XP002067998 see page 2324, scheme 2 -----	1-8, 11-17